



Because we care for your birds

# **PANTACOX**

# WATER SOLUBLE LIQUID 100 ML

## TREATMENT OF COCCIDIOSIS

**Coccidiosis** is an infectious disease caused by protozoa (small single-celled parasites). It develops in the host's *intestinal cells* and damages the intestinal wall. Digestion is then severely impaired and birds lose weight. Young birds are very susceptible!

**Atoxoplasmosis** is also a specific form of coccidiosis. This parasite enters the bloodstream during its cycle and can settle in organs such as the liver, spleen, lungs and brain. Birds develop thickened livers, bloody spots on the intestines, wheezing and sometimes loss of balance.

**Pantacox** intervenes in the coccidiosis cycle and kills coccidia. It is therefore a coccidiocide. It has no negative effects on fertilisation, laying, growth or hatching outcomes.

# PANTACOX—COCCIDIOSIS TREATMENT CARE BRIDGS AND PRESENTE COCCIDIOSIS TREATMENT COCCIDIOSIS TREATMENT COME MANY AND PRINTER OF M

### **INDICATIONS**

- Preventive treatment before the start of breeding.
- Treatment of coccidiosis infection.
- Control of atoxoplasmosis (Lankesterella).
- Preventive protection during the moulting period.

## **DOSAGE AND ADMINISTRATION**

Oral administration via drinking water

 In case of coccidiosis: 2 to 3 ml per litre of drinking water for 2 to 3 consecutive days. If necessary, repeat the treatment after two weeks.

 In case of atoxoplasmosis: 2 to 3 ml per litre of drinking water for 2 consecutive days, then continue treatment with ESB PLUS.

Change the medicinal water daily and provide it as the only source of drinking water during treatment. Additional vitamins can be given during rest periods.

Each pack contains a 1 ml syringe.



Ordering through:

WWW.PANTEX-COUTTEEL.COM